

- paleoceanography, and depositional processes (abstract): AAPG Mid-Continent Meeting Official Program, p. 21.
- Conant, L.C., and V.E. Swanson, 1961, Chattanooga Shale and related rocks of central Tennessee and nearby areas: U.S. Geological Survey Professional Paper 357, 91 p.
- Cook, T.D., and A.W. Bally, eds., 1975, Stratigraphic atlas of North and Central America: Princeton University Press, 272 p.
- Cooper, C.L., 1931a, Conodonts from the Arkansas novaculite, Woodford Formation, Ohio Shale, and Sunbury Shale: *Journal of Paleontology*, v. 5, p. 143-151.
- Cooper, C.L., 1931b, New conodonts from the Woodford Formation of Oklahoma: *Journal of Paleontology*, v. 5, p. 230-243.
- Cooper, C.L., 1932, A crustacean fauna from the Woodford Formation of Oklahoma: *Journal of Paleontology*, v. 6, p. 346-352.
- Cooper, C.L., 1935, Conodonts from the upper and middle Arkansas novaculite, Mississippian, at Caddo Gap, Arkansas: *Journal of Paleontology*, v. 9, p. 307-315.
- Craig, W.W., O. Wise, and J.D. McFarland, III, 1984, A guidebook to the post-St. Peter Ordovician and the Silurian and Devonian rocks of north-central Arkansas: Arkansas Geological Commission, 49 p. (Chattanooga Shale, p. 21-23)
- Crick, R.E., B.B. Ellwood, D.J. Over, R. Feist, and C. Girard, 2001, Magnetostratigraphic susceptibility of the Frasnian-Famennian boundary (Upper Devonian) in southern Oklahoma and its relationship to the type area in southern France, *in* K.S. Johnson, ed., Silurian, Devonian, and Mississippian geology and petroleum in the southern Midcontinent, 1999 symposium: OGS Circular 105, p. 71-82.
- Crossey, L.J., E.S. Hagen, R.C. Surdam, and T.W. Papoint, 1986, Correlation of organic parameters derived from elemental analysis and programmed pyrolysis of kerogen, *in* D.L. Gautier, ed., Roles of organic matter in sediment diagenesis: SEPM Special Publication 38, p. 35-45.
- Cruse, A., S.T. Paxton, and M. Aufill, 2009, Constraints on sedimentation rates of the Woodford Shale from modeling of uranium diffusion (abstract): AAPG Mid-Continent Meeting Official Program, p. 21.
- Curtis, M.E., B.J. Cardott, C.H. Sondergeld, and C.S. Rai, 2012, Development of organic porosity in the Woodford Shale with increasing thermal maturity: *International Journal of Coal Geology*, v. 103, p. 26-31.
- Damsté, J.S.S., T.I. Eglinton, J.W. de Leeuw, and P.A. Schenck, 1989, Organic sulphur in macromolecular sedimentary organic matter: I. Structure and origin of sulphur-containing moieties in kerogen, asphaltenes and coal as revealed by flash pyrolysis: *Geochimica et Cosmochimica Acta*, v. 53, p. 873-889. (Woodford kerogen)
- Davis, H.G., 1985, Wrenching and oil migration, Mervine field, Kay County, Oklahoma [abstract]: AAPG Bulletin, v. 69, p. 248.
- Davis, H.G., 1994, Petroleum geology of the Woodford Formation, Anadarko basin, Oklahoma and Texas [abstract]: AAPG Annual Convention, Official Program, v. 3, p. 132.